



Tobacco Use and Cessation in Adolescents: A Review

Merazga Batoul 

¹ Clinical psychology laboratory, University mohamed lamine debaghine (Algeria),
ba.merazga@univ-setif2.dz
( 0009-0009-2225-6430)

Received: 03/07/2025

Accepted: 23/05/2026

Abstract:

This review aims to explore the global prevalence of smoking among adolescents and examine the underlying factors contributing to its initiation and persistence. Key influences such as genetic predisposition, family environment, peer dynamics, and media exposure are discussed in light of their impact on youth behavior.

The paper highlights the serious physical, psychological, and social consequences of tobacco use and evaluates evidence-based cessation strategies, including behavioral therapies, pharmacological tools, and integrative approaches. Emphasis is placed on the importance of early intervention, psychological support, and culturally tailored policies to promote lasting change among adolescents.

Keywords: Adolescent Smoking; Health Psychology; Nicotine Addiction; Prevention Strategies; Smoking Cessation; Tobacco Use

1. INTRODUCTION

Tobacco use remains one of the most preventable yet persistent contributors to global illness and premature death. Nicotine—the addictive component in tobacco—is classified as a psychoactive substance and is central to what is now clinically recognized as Tobacco Use Disorder (TUD) in the DSM-5 and Nicotine Dependence under ICD-10 criteria (American Psychiatric Association, 2013; WHO, 2020). Despite numerous awareness initiatives, tobacco continues to be widely available and socially accepted in many environments, particularly among teenagers who are highly susceptible to peer pressure, familial behaviors, and targeted advertising campaigns (Mermelstein, 1999; Dai et al., 2022).

The Global Burden of Disease study estimates that around 1.3 billion people worldwide use tobacco, with a staggering 80% residing in low- and middle-income nations (GBD, 2019). Each year, tobacco use results in over 8 million deaths globally, including roughly 1.3 million from exposure to secondhand smoke (WHO, 2023). Most smokers begin before turning 18, and those who start early often face greater difficulty quitting later in life (Lasser et al., 2000; Piper et al., 2022). In addition, adolescent smokers are more vulnerable to developing mental health challenges, such as depression, anxiety, and substance misuse (Chaiton et al., 2009; Hall et al., 2016).

During adolescence, the brain is still undergoing major developmental changes, making it especially sensitive to nicotine's neurochemical effects. This vulnerability increases the likelihood of forming long-term dependencies and experiencing cognitive impairments (Goriounova & Mansvelder, 2012). Furthermore, adolescent smoking has been linked to academic struggles, disturbed sleep patterns, and overall lower levels of life satisfaction (Leung et al., 2017; West, 2017). Starting to smoke at a young age also elevates the lifetime risk for chronic conditions such as coronary artery disease, stroke, chronic obstructive pulmonary disease (COPD), and a variety of cancers, particularly those affecting the lungs, pancreas, and esophagus (Chandrupatla et al., 2017; Rigotti et al., 2022).

The rapid expansion of digital platforms has added new challenges to tobacco control. Adolescents are increasingly exposed to pro-smoking messages on social media, which often glamorize or normalize tobacco use (Donaldson et al., 2022). Meanwhile, the popularity of vaping and e-cigarettes continues to grow, especially among youth, despite growing evidence linking these products to health risks and traditional cigarette use (Wang et al., 2021; O'Brien et al., 2021).

Given these concerning trends, this review seeks to offer a comprehensive analysis of adolescent tobacco use. It explores the global prevalence, risk factors, and health consequences associated with youth smoking. The paper also critically examines existing behavioral and medical cessation strategies from a health psychology perspective, advocating for integrated approaches that address not only individual behaviors but also the broader social and cultural context in which tobacco use occurs.

2. Global Prevalence of Tobacco Use Among Adolescents

The initiation of tobacco use often takes place during the teenage years, with research indicating that up to 90% of adult smokers had their first experience with cigarettes between the ages of 15 and 25 (Warner, 2019). Although significant reductions in youth smoking have been recorded in many high-income countries, the global picture remains troubling. Tobacco use continues to persist in numerous regions, especially in low- and middle-income countries, where enforcement of tobacco regulations is often weak (Dai et al., 2022).

According to the World Health Organization (WHO), as of 2020, about 32.6% of

men and 6.5% of women worldwide were active smokers, totaling more than 1.18 billion users globally (WHO, 2021). The sharpest declines in smoking rates have occurred in countries with strong public health infrastructure, high educational levels, and strict tobacco control policies. Nations like Denmark, Brazil, Nepal, and the Netherlands have seen smoking prevalence drop by more than 40% since 1990 (Gakidou et al., 2020).

In contrast, progress in reducing smoking rates has been slower in parts of Asia and Africa. Countries such as China and Indonesia still report alarmingly high smoking prevalence among adult males, often exceeding 50% (GBD, 2021; Dai et al., 2022). A recent systematic review by Jha and colleagues (2023) indicates that adolescent smoking in regions like Sub-Saharan Africa and Southeast Asia remains steady or is even on the rise—largely due to pervasive tobacco marketing, cultural acceptance of smoking, and minimal enforcement of age-related sales restrictions.

Despite widespread awareness of smoking's dangers, many adolescents who start smoking struggle to quit. Around 70% of smokers report a desire to stop, yet most relapse within their first few attempts. On average, individuals make between five and seven serious attempts before achieving long-term abstinence (Rigotti et al., 2022). These numbers reflect both the addictive nature of nicotine and the challenges associated with sustaining behavior change.

Recent analyses from the Cochrane Tobacco Addiction Group underscore the importance of early intervention. School-based prevention programs, when implemented effectively, can delay the age of smoking initiation and reduce smoking rates among students (Thomas et al., 2021). Nonetheless, tobacco use among adolescents remains a major global health concern, demanding ongoing attention through culturally sensitive and region-specific strategies.

These findings remind us that while progress is possible, it remains uneven and fragile. The persistence of adolescent smoking, particularly in underserved regions, reflects deeper systemic issues—such as socioeconomic inequality, weak governance, and aggressive marketing by tobacco companies. Moreover, prevalence rates may not always reflect the true scope of the problem, especially in underdeveloped societies where data collection is limited or unreliable. In such contexts, smoking—particularly among adolescents—often goes underreported or socially normalized. This reinforces the urgent need for culturally sensitive, evidence-based interventions that not only target behavior but also confront the structural roots of tobacco use as a public health crisis.

3. Risk Factors Contributing to Adolescent Smoking Behavior

Adolescent smoking is rarely the result of a single cause. Instead, it stems from a complex web of interacting influences, including biological, psychological, social, and environmental factors. Research consistently shows that multiple elements contribute to the decision to start—and continue—smoking during adolescence.

-Gender Differences: Studies have found that adolescent males are more likely to smoke than females, particularly in cultural contexts where male smoking is socially accepted or encouraged (Öncel et al., 2011).

-Parental Influence: Adolescents with one or both parents who smoke are significantly more likely to take up smoking themselves. Parental modeling and household smoking norms play a critical role in shaping behavior (Hesketh et al., 2001).

-Family Environment: Teens from families experiencing divorce, parental death, or other disruptions tend to have higher rates of tobacco use, potentially due to emotional distress or reduced supervision (Donaldson et al., 2022).

-Socioeconomic Status (SES): Both ends of the SES spectrum are linked to smoking, though the underlying causes differ. For instance, low-income households may foster smoking due to stress, while higher-income youth may be influenced by social trends or peer dynamics (Ehsani-Chimeh et al., 2020).

-Parental Education: Adolescents whose parents have lower levels of education are more likely to smoke. This may reflect differences in health literacy and the ability to convey anti-smoking attitudes effectively (Öncel et al., 2011).

-Peer Pressure: The behavior of close friends is one of the strongest predictors of adolescent smoking. Teens who have peers who smoke are far more likely to try smoking themselves (Chung & Joung, 2014; Leventhal et al., 2020).

-School Engagement: Poor academic performance, disengagement from school, and lack of involvement in extracurricular activities are all associated with a higher likelihood of smoking during adolescence (West, 2017).

-Mental Health Issues: Conditions such as depression, anxiety, and conduct disorders significantly increase the risk of smoking initiation and maintenance among teenagers (Lasser et al., 2000; Hall et al., 2016).

-Substance Use: Smoking often co-occurs with other high-risk behaviors, such as alcohol or drug use, creating a reinforcing cycle of risk (Tariq & Gupta, 2020).

-Media Exposure: Adolescents exposed to smoking in movies, on television, and on social media platforms are more likely to develop favorable attitudes toward tobacco and to perceive it as a socially acceptable behavior (Donaldson et al., 2022).

-Accessibility: In many countries, weak regulations allow teenagers to obtain cigarettes easily, making experimentation and regular use more likely (Ehsani-Chimeh et al., 2020).

-Perception of Risk: When adolescents underestimate the health risks of smoking, they become more susceptible to environmental cues and social pressure (Wang et al., 2021). Moreover, the influence of social media personalities and online trends cannot be overlooked. Platforms such as TikTok, Instagram, and YouTube often present smoking as fashionable or rebellious, which can shape adolescents' beliefs and increase their likelihood of trying tobacco (Majmundar et al., 2023).

Overall, these diverse risk factors highlight the need for multilayered prevention efforts. Effective strategies should address individual, familial, school, and societal influences simultaneously to curb adolescent smoking and prevent long-term addiction.

4. Health Consequences of Smoking

The health implications of tobacco use are both extensive and severe, affecting not only those who smoke directly but also individuals exposed to secondhand smoke. While nicotine is the substance that drives addiction, the true harm comes from the toxic chemicals present in tobacco smoke—compounds like arsenic, formaldehyde, benzene, and carbon monoxide, all of which contribute to widespread physiological damage (West, 2017; WHO, 2023).

Among the major health outcomes associated with smoking are:

Respiratory Diseases: Cigarette smoke is a primary cause of chronic obstructive pulmonary disease (COPD), which includes conditions such as chronic bronchitis and emphysema. It is also linked to other serious lung diseases, including pulmonary fibrosis (Rigotti et al., 2022).

Cardiovascular Complications: Smoking accelerates the buildup of plaque in arteries, a process known as atherosclerosis, which significantly increases the risk of heart attacks, strokes, high blood pressure, and peripheral vascular disease (Nicolosi et al., 2003).

Cancer: Tobacco use is responsible for more than a dozen types of cancer, with the lungs, throat, pancreas, kidneys, bladder, and esophagus among the most commonly affected organs (Chandrupatla et al., 2017; GBD, 2019).

Reproductive and Sexual Health: In males, smoking is associated with erectile dysfunction, while in women, it increases the risk of complications during pregnancy, such as preterm birth and low birth weight (Warner & Mendez, 2019).

Immune Function and Recovery: Smokers are more prone to infections and

autoimmune disorders, and they generally experience slower healing times after injury or surgery (West, 2017).

COVID-19 Outcomes: Evidence suggests that smokers are more likely to experience severe COVID-19 symptoms and are at greater risk of death from the virus compared to nonsmokers (Patanavanich & Glantz, 2020).

Beyond physical health, smoking has been consistently linked to adverse psychological effects. Adolescents who smoke are more likely to report symptoms of depression, anxiety, and suicidal thoughts than their nonsmoking peers (Hall et al., 2016; Chaiton et al., 2009).

There are also significant financial and societal costs associated with tobacco use. These include not only direct healthcare expenses but also indirect burdens such as loss of productivity, premature death, and money spent on tobacco products (WHO, 2021). Altogether, the wide-ranging consequences of smoking emphasize the urgent need for effective prevention and cessation efforts—particularly among young people, whose developing bodies and brains are especially vulnerable to tobacco’s harmful effects.

5. Evidence-Based Interventions for Smoking Cessation

Although nicotine addiction presents a considerable barrier to quitting, a range of scientifically supported interventions are available to assist adolescents in overcoming tobacco dependence. These strategies fall into three broad categories: behavioral, pharmacological, and hybrid approaches.

5.1. Motivational Interviewing (MI)

Motivational Interviewing is a collaborative, client-centered counseling method designed to strengthen personal motivation for change. By helping individuals resolve ambivalence about quitting, MI encourages behavior modification in a non-confrontational manner (Miller & Rollnick, 2013). While outcomes can vary, MI has shown promise in adolescent populations, particularly among those not yet fully committed to quitting (Lindson et al., 2019).

5.2. Acceptance and Commitment Therapy (ACT)

ACT is a third-wave cognitive-behavioral approach that focuses on accepting cravings and unhelpful thoughts without acting on them. Rather than trying to eliminate urges, ACT guides individuals to align their actions with personal values—such as health or family—while defusing from smoking-related thoughts (Hayes et al., 2013). Controlled trials suggest ACT can outperform traditional behavioral therapies in helping individuals reduce smoking behavior (Gifford et al., 2011).

5.3. Pharmacological Treatments

Several medications have been developed to aid smoking cessation by addressing nicotine dependence and withdrawal symptoms. Common pharmacological options include:

- **Nicotine Replacement Therapy (NRT):** Available in various forms such as patches, gums, and lozenges, NRT delivers controlled doses of nicotine to ease withdrawal and reduce cravings.
- **Varenicline (Chantix):** A partial agonist that targets nicotinic receptors, varenicline has demonstrated high success rates in promoting abstinence.
- **Bupropion (Zyban):** Originally developed as an antidepressant, bupropion also helps reduce nicotine cravings and withdrawal symptoms.

Meta-analyses show that combining behavioral support with medication substantially improves quit rates. In fact, research indicates that the success rate nearly doubles—from about 8.6% with minimal support to over 15% with combined intervention (Rigotti et al., 2022).

Recent results from the EAGLES trial also show that varenicline leads the way in single-agent effectiveness, with a 21.8% quit rate after six months, followed by bupropion at 16.2%, and nicotine patches at 15.7% (Rigotti et al., 2023).

Another emerging option is **cytisinicline**, a plant-based compound that has recently attracted attention as a potential cessation aid. A U.S.-based clinical trial involving over 800 adult smokers found cytisinicline to be more effective than placebo and well-tolerated, with fewer side effects than some existing medications (Rigotti et al., 2023).

5.4. Electronic Cigarettes and Vaping Devices

E-cigarettes function by vaporizing a nicotine-containing liquid, offering an alternative to combustible tobacco. While they are often perceived as less harmful, the long-term effects of vaping remain uncertain (Lindson et al., 2024). Some public health experts argue that e-cigarettes can serve as transitional tools for adult smokers who are trying to quit, particularly when used under medical supervision (Wang et al., 2021). However, concerns remain regarding dual use and the possibility that adolescents who vape may progress to conventional smoking (O'Brien et al., 2021).

Despite the well-documented and wide-ranging health risks of tobacco use—from chronic diseases and cancer to psychological distress and socioeconomic burden—many smokers continue the habit, often for years or decades. This persistence highlights the complex and addictive nature of smoking, which extends beyond physiological dependence on nicotine to include deep-seated behavioral and psychological components. Many smokers are aware of the dangers but struggle to quit due to emotional triggers, habitual routines, and perceived benefits, such as stress relief. This underscores the need for professional intervention and for treating tobacco use as a form of behavioral addiction. In the field of health psychology, one of the most widely applied frameworks for understanding and supporting smoking cessation is the Transtheoretical Model of Behavior Change (TTM) (Dadras, 2024). This model emphasizes the importance of guiding individuals through progressive stages—from precontemplation to maintenance—offering tailored interventions that correspond to the smoker's readiness to change. Integrating psychological support with medical strategies increases the chances of successful, long-term cessation.

6. Transtheoretical Model of Behavior Change

The Stages of Change model, proposed by James Prochaska and Carlo DiClemente in the 1980s (Prochaska, 2008), was developed to better understand how individuals transition out of unhealthy behaviors. Originally based on research into smoking cessation, the model has since been applied across a variety of health-related behaviors, including substance use, dietary change, and physical activity. The framework outlines six key stages: precontemplation, contemplation, preparation, action, maintenance, and sometimes relapse.

In the *precontemplation stage*, the smoker typically does not recognize their behavior as harmful. They may feel defensive, minimize the risks, or express doubt about their

ability to quit. Individuals in this stage often avoid health warnings or downplay symptoms they may be experiencing. As they move into the *contemplation stage*, they begin to reflect more seriously on the health and social consequences of smoking. There is internal conflict: they may want to quit but feel uncertain or afraid of change. This stage can last months or even years.

The preparation stage marks a more active commitment to change. Here, the individual may start reducing cigarette intake, research cessation methods, or seek support from healthcare professionals. They often set a quit date and mentally prepare for the challenges ahead. *The action stage* involves a full commitment to quitting. The smoker stops using tobacco and replaces the habit with healthier behaviors. This stage requires coping strategies, such as avoiding triggers, managing stress, and staying socially supported.

Once in *the maintenance stage*, the individual works to sustain the change over time. They may experience occasional cravings or emotional stress, but they have developed tools to prevent relapse. However, *relapse* can occur at any stage and is not a sign of failure but rather part of the learning process. Returning to earlier stages allows individuals to strengthen their motivation and re-strategize for future attempts.

From a human perspective, it's important to recognize that the resistance to change often increases with the duration of smoking and with age. Individuals who have smoked for decades may feel that quitting is nearly impossible or unnecessary, especially if they've attempted and failed multiple times. Long-term smokers often link tobacco use to their identity, daily routine, or emotional regulation, making change not only a physical challenge but also a psychological upheaval. Feelings of fear, loss, and even grief can arise when letting go of a habit that has served as a coping mechanism for years.

Moreover, older smokers may be less exposed to modern anti-smoking campaigns or feel excluded from youth-centered prevention efforts. They might also face health issues that reduce their confidence in their ability to change. Therefore, interventions should be tailored not only to the individual's stage of change but also to their personal history, emotional readiness, age, and belief systems. Empathy, patience, and a nonjudgmental attitude are essential. The path to quitting is rarely linear, but with consistent support and individualized care, meaningful change remains possible—at any age, and at any stage.

7. Benefits of Smoking Cessation

Quitting smoking delivers a wide range of health, psychological, and financial benefits—many of which begin within hours or days after cessation and continue to improve over time. These advantages are not limited to the individual smoker but also extend to those around them, particularly vulnerable populations like children and pregnant women (Donaldson et al., 2022).

-Cardiovascular Health: One of the most immediate benefits of quitting is improved heart health. Within the first one to two years of cessation, the risk of developing coronary heart disease drops significantly and continues to decline with each smoke-free year (Rigotti et al., 2022).

-Reduced Cancer Risk: Smoking cessation greatly decreases the risk of developing various cancers, including those affecting the lungs, bladder, pancreas, mouth, and throat. The risk reduction strengthens with time, especially among those who quit at younger ages (GBD, 2019).

-Respiratory Improvement: Lung function begins to recover within weeks after quitting. Former smokers often report less coughing, improved breathing, and reduced shortness of breath (West, 2017).

-Enhanced Immune Function and Fertility: Stopping smoking helps the immune system recover and lowers the chances of developing infections. It also improves fertility, particularly for women in their reproductive years (Warner & Mendez, 2019).

Lower Stroke and Vascular Disease Risk: Smoking cessation reduces the likelihood of stroke and other cerebrovascular complications by improving blood vessel function and circulation (Nicolosi et al., 2003).

-Better Mental Health: Beyond physical recovery, many individuals experience improved psychological well-being. Studies show significant reductions in symptoms of depression, anxiety, and stress within the first year of quitting, particularly in young adults and adolescents (Hall et al., 2016).

-Financial Gains: Over time, the money saved from not purchasing tobacco products can be substantial. For adolescents and young adults, this may also redirect spending toward healthier lifestyle choices or educational opportunities.

-Protection of Others: Quitting smoking also eliminates the harm posed by secondhand smoke. Children, pregnant women, and individuals with chronic illnesses benefit most from reduced exposure in shared environments.

Ultimately, the advantages of quitting smoking are profound, cumulative, and well-supported by decades of medical research. These outcomes reinforce the urgency of early intervention and the promotion of accessible cessation support—especially for adolescents, who stand to gain the most from stopping tobacco use early in life.

8. Recommendations and Future Directions

Despite growing awareness of the dangers associated with tobacco use, smoking among adolescents remains a persistent public health challenge. To combat this issue effectively, coordinated, multi-level strategies must be implemented. The following recommendations highlight key areas for action:

- **Strengthen Early Prevention Initiatives**

Schools and community programs should prioritize early education on the harms of tobacco, emphasizing skill-building, peer resistance, and awareness of industry manipulation tactics. Evidence shows that early intervention can delay or prevent smoking initiation.

- **Tighten Regulations on Tobacco Marketing**

Particular attention should be paid to controlling tobacco-related content on social media and digital platforms, which frequently target younger audiences through influencers, music videos, and trending challenges.

3. Involve Families in Intervention Efforts

Parents play a vital role in shaping adolescents' attitudes toward smoking. Family-based programs that promote smoke-free homes, positive role modeling, and open communication can reduce the likelihood of youth tobacco use.

- **Improve Access to Cessation Resources**

Behavioral therapies and pharmacological aids should be made widely available, particularly in underserved communities. Health systems and schools must collaborate to ensure these services are affordable, youth-friendly, and culturally appropriate.

- **Expand Digital Support Tools**

The development and deployment of mobile health (mHealth) tools—such as apps, text-based coaching, and online counseling—can provide scalable and discreet support for teens seeking to quit.

- **Prohibit Flavored Tobacco Products**

Flavored cigarettes and vape liquids disproportionately appeal to young people. Policies that restrict or ban these products could significantly reduce initiation rates (CDC, 2022).

- **Integrate Smoking Cessation into Mental Health Services**

Given the strong link between tobacco use and mental health, therapists, school counselors, and pediatricians should be trained to incorporate smoking assessments and cessation strategies into routine care.

- **Support Research on Emerging Interventions**

New cessation aids like cytisinicline and electronic nicotine delivery systems require further evaluation, particularly in adolescent populations. Longitudinal studies can provide deeper insights into long-term safety and effectiveness.

- **Promote Culturally Tailored Interventions**

Tobacco control programs should be adapted to local cultural norms, especially in communities where smoking is normalized or tied to gender identity or social status. Engaging community leaders and local influencers can improve reach and effectiveness.

By addressing the root causes of adolescent smoking and tailoring responses to specific sociocultural environments, these recommendations align with a health psychology framework that values context, behavior, and long-term change. Collective action from policymakers, educators, families, and healthcare providers will be essential to protect the health of future generations.

With global cooperation and targeted intervention, a future free from tobacco-related harm among youth is within reach.

9. Conclusion

Adolescent tobacco use remains a complex and deeply rooted public health concern, driven by a web of biological, psychological, social, and cultural influences. The addictive properties of nicotine, combined with the powerful sway of peer dynamics, media exposure, and familial patterns, make it especially difficult for young people to quit once they start. Despite widespread awareness of the risks, smoking continues to claim millions of lives each year—and many of those lives are shaped by habits formed in adolescence.

This review highlights that while the challenge is formidable, there are promising pathways forward. A growing body of evidence supports the effectiveness of behavioral and pharmacological interventions tailored to youth, including motivational interviewing, acceptance-based therapies, nicotine replacement strategies, and emerging tools like cytisinicline and regulated e-cigarette use. When applied within a broader framework that includes education, regulation, family involvement, and culturally

responsive outreach, these interventions can significantly reduce smoking initiation and support long-term cessation.

Health psychology provides a valuable lens for understanding how personal, social, and environmental factors shape behavior—and how those same pathways can be used to foster change. As the global landscape of tobacco use continues to evolve, integrating established methods with innovative solutions will be essential. Prioritizing prevention, enhancing access to care, and investing in youth-centered approaches will not only reduce tobacco-related harm but also contribute to the overall health and well-being of future generations.

Referrals and references:

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: Author.
- Chandrupatla, S. G., Tavares, M., & Natto, Z. S. (2017). Tobacco use and effects of professional advice on smoking cessation among youth in India. *Asian Pacific Journal of Cancer Prevention*, 18(7), 1861–1867. <https://doi.org/10.22034/APJCP.2017.18.7.1861>
- Chaiton, M., Cohen, J. E., O'Loughlin, J., & Rehm, J. (2009). A systematic review of longitudinal studies on the association between depression and smoking in adolescents. *BMC Public Health*, 9(1), 1–11. <https://doi.org/10.1186/1471-2458-9-356>
- Chung, S. S., & Joung, K. H. (2014). Risk factors for smoking behaviors among adolescents. *The Journal of School Nursing*, 30(4), 262–271. <https://doi.org/10.1177/1059840513505222>
- CDC. (2022). *Smoking & tobacco use: Fast facts*. Centers for Disease Control and Prevention. https://www.cdc.gov/tobacco/data_statistics/fact_sheets
- Dadras, O. (2024). Predictor of smoking cessation among school-going adolescents in Indonesia: a secondary analysis based on the transtheoretical model of behavioral change. *Frontiers in Psychiatry*, 15, 1374731.
- Dai, X., Gakidou, E., & Lopez, A. D. (2022). Evolution of the global smoking epidemic over the past half century: Strengthening the evidence base for policy action. *Tobacco Control*, 31(2), 129–137. <https://doi.org/10.1136/tobaccocontrol-2021-056535>
- Donaldson, S. I., Dormanesh, A., Perez, C., Majmundar, A., & Allem, J. P. (2022). Association between exposure to tobacco content on social media and tobacco use: A systematic review and meta-analysis. *JAMA Pediatrics*, 176(5), 487–495. <https://doi.org/10.1001/jamapediatrics.2021.6415>
- Ehsani-Chimeh, E., Sajadi, H. S., Behzadifar, M., et al. (2020). Current and former smokers among adolescents aged 12–17 years in Iran: A systematic review and meta-analysis. *BMC Public Health*, 20(1), 1–15. <https://doi.org/10.1186/s12889-020-8255-2>
- GBD 2019 Tobacco Collaborators. (2021). Prevalence of smoking and attributable disease burden in 204 countries, 1990–2019. *The Lancet*, 397(10292), 2337–2360. [https://doi.org/10.1016/S0140-6736\(21\)01169-7](https://doi.org/10.1016/S0140-6736(21)01169-7)
- Gifford, E. V., Kohlenberg, B. S., Hayes, S. C., et al. (2011). Acceptance-based treatment for smoking cessation. *Behavior Therapy*, 42(4), 689–700. <https://doi.org/10.1016/j.beth.2011.01.003>
- Goriounova, N. A., & Mansvelder, H. D. (2012). Short- and long-term consequences of nicotine exposure during adolescence for prefrontal cortex neuronal network function. *Cold Spring Harbor Perspectives in Medicine*, 2(12), a012120. <https://doi.org/10.1101/cshperspect.a012120>
- Hall, S. M., Prochaska, J. J., & Dilley, J. W. (2016). The relationship between smoking and mental illness: Revisited. *Annual Review of Clinical Psychology*, 12, 409–431. <https://doi.org/10.1146/annurev-clinpsy-021815-093734>
- Hayes, S. C., Levin, M. E., Plumb-Villardaga, J., Villatte, J. L., & Pistorello, J. (2013). Acceptance and commitment therapy: Progress of a distinctive model. *Behavior Therapy*, 44(2), 180–198. <https://doi.org/10.1016/j.beth.2009.08.002>
- Hesketh, T., Ding, Q. J., & Tomkins, A. (2001). Smoking among youths in China. *American Journal of Public Health*, 91(10), 1653–1655. <https://doi.org/10.2105/AJPH.91.10.1653>
- Jha, P., MacLennan, M., Chaloupka, F. J., et al. (2023). Global tobacco economics: Challenges and prospects. *The Lancet Global Health*, 11(4), e537–e548. [https://doi.org/10.1016/S2214-109X\(23\)00007-1](https://doi.org/10.1016/S2214-109X(23)00007-1)
- Lasser, K., Boyd, J. W., Woolhandler, S., Himmelstein, D. U., McCormick, D., & Bor, D. H. (2000). Smoking and mental illness: A population-based prevalence study. *JAMA*, 284(20), 2606–2610. <https://doi.org/10.1001/jama.284.20.2606>

- Leung, L. T., Toumbourou, J. W., & Hemphill, S. A. (2017). The effect of peer influence and school engagement on adolescent smoking. *Addictive Behaviors*, 65, 20–25. <https://doi.org/10.1016/j.addbeh.2016.10.009>
- Lindson, N., Butler, A. R., McRobbie, H., et al. (2024). Electronic cigarettes for smoking cessation. *Cochrane Database of Systematic Reviews*, 1, CD015554. <https://doi.org/10.1002/14651858.CD015554.pub2>
- Lindson, N., Thompson, T. P., Ferrey, A., Lambert, J. D., & Aveyard, P. (2019). Motivational interviewing for smoking cessation. *Cochrane Database of Systematic Reviews*, (7), CD006936. <https://doi.org/10.1002/14651858.CD006936.pub4>
- Majmundar, A., Allem, J. P., & Unger, J. B. (2023). The impact of influencer marketing on youth smoking behavior. *Health Psychology Review*, 17(1), 20–36. <https://doi.org/10.1080/17437199.2022.2148375>
- Mermelstein, R. (1999). Ethnicity, gender and risk factors for smoking initiation: An overview. *Nicotine & Tobacco Research*, 1(Suppl_2), S39–S43. <https://doi.org/10.1080/14622299050011791>
- Miller, W. R., & Rollnick, S. (2013). *Motivational interviewing: Helping people change* (3rd ed.). New York: Guilford Press.
- Nicolosi, A., Moreira Jr, E. D., Shirai, M., Tambi, M. I., & Glasser, D. B. (2003). Epidemiology of erectile dysfunction in four countries. *Urology*, 61(1), 201–206. [https://doi.org/10.1016/S0090-4295\(02\)02194-7](https://doi.org/10.1016/S0090-4295(02)02194-7)
- O'Brien, D., Long, J., Quigley, J., et al. (2021). E-cigarettes and tobacco smoking initiation in adolescents: A systematic review and meta-analysis. *BMC Public Health*, 21(1), 1–10. <https://doi.org/10.1186/s12889-021-10655-8>
- Öncel, S. Y., Gebizlioğlu, Ö. L., & Alioğlu, F. A. (2011). Risk factors for smoking behavior among university students. *Turkish Journal of Medical Sciences*, 41(6), 1071–1080. <https://doi.org/10.3906/sag-1009-1122>
- Patanavanich, R., & Glantz, S. A. (2020). Smoking and COVID-19: A meta-analysis. *Nicotine & Tobacco Research*, 22(9), 1653–1656. <https://doi.org/10.1093/ntr/ntaa082>
- Piper, M. E., Fowler, C. D., Serena, T., et al. (2022). Tobacco and nicotine use. *Nature Reviews Disease Primers*, 8(1), 1–21. <https://doi.org/10.1038/s41572-022-00352-8>
- Prochaska, J. O. (2008). Decision making in the transtheoretical model of behavior change. *Medical decision making*, 28(6), 845–849.
- Rigotti, N. A., Benowitz, N. L., Prochaska, J., et al. (2023). Cytisinicline for smoking cessation: A randomized clinical trial. *JAMA*, 330(2), 152–160. <https://doi.org/10.1001/jama.2023.10042>
- Rigotti, N. A., Kruse, G. R., Livingstone-Banks, J., & Hartmann-Boyce, J. (2022). Treatment of tobacco smoking: A review. *JAMA*, 327(6), 566–577. <https://doi.org/10.1001/jama.2022.1923>
- Tariq, N., & Gupta, V. (2020). High risk behaviors. In *StatPearls*. StatPearls Publishing. <https://www.ncbi.nlm.nih.gov/books/NBK560756/>
- Thomas, R. E., McLellan, J., & Perera, R. (2021). School-based programmes for preventing smoking. *Cochrane Database of Systematic Reviews*, 10, CD001293. <https://doi.org/10.1002/14651858.CD001293.pub5>
- Wang, R. J., Bhadriraju, S., & Glantz, S. A. (2021). E-cigarette use and adult smoking cessation: A meta-analysis. *American Journal of Public Health*, 111(2), 230–246. <https://doi.org/10.2105/AJPH.2020.306087>
- Warner, K. E. (2019). How to think—not feel—about tobacco harm reduction. *Nicotine & Tobacco Research*, 21(10), 1299–1309. <https://doi.org/10.1093/ntr/nty084>
- Warner, K. E., & Mendez, D. (2019). E-cigarettes: Comparing the possible risks of increasing smoking initiation with the potential benefits of increasing smoking cessation. *Nicotine & Tobacco Research*, 21(1), 41–47. <https://doi.org/10.1093/ntr/ntx166>
- West, R. (2017). Tobacco smoking: Health impact, prevalence, correlates and interventions. *Psychology & Health*, 32(8), 1018–1036. <https://doi.org/10.1080/08870446.2017.1325890>
- World Health Organization (WHO). (2020). *International Classification of Diseases (ICD-10)*. <https://icd.who.int/>
- World Health Organization (WHO). (2021). *Global report on trends in prevalence of tobacco use 2000–2025* (3rd ed.). <https://www.who.int/publications/i/item/9789240039322>
- World Health Organization (WHO). (2023). *Tobacco: Key facts*. <https://www.who.int/news-room/fact-sheets/detail/tobacco>