

A NUMERICAL METHOD FOR SOLVING A TWO-DIMENSIONAL DIFFUSION EQUATION WITH NON LOCAL BOUNDARY CONDITIONS

A. CHENIGUEL

Department of mathematics and computer science, Faculty of sciences
Larbi ben mhidi University, Oum-Elbouaghi, Algeria.

Reçu le 10/10/2010 – Accepté le 27/05/2011

Résumé

Dans ce travail on résout quelques problèmes aux limites avec des conditions aux limites non locales dans le cas bidimensionnel . En utilisant la méthode de décomposition, la solution analytique est calculée sous la forme d'une série entière, les différents termes de cette série sont facile à calculer. Les résultats obtenus sont tous exactes d'où l'avantage de cette méthode par rapport aux méthodes classiques telles que les méthodes des différences finis, éléments finis, spectrale....

Mots clés: Monodimensionnelles, mouvement de l'atmosphère, transition de phase gaz liquide

Abstract

This paper is devoted to the decomposition method which is applied to solve problems with non local boundary conditions. The analytic solution of the problem is calculated in a series form with easily computable components. The comparison of the methodology with some known techniques shows that the present approach is powerful, efficient and reliable.

Keywords: decomposition method, non local boundary conditions, partial differential equations, Analytic solution

ملخص

نقترح في هذا البحث حلا تقريبا لبعض مسائل القيمة الحدية بشروط حدية غير موضعية و لذلك نستعمل طريقة التفكيك و ذلك بفرض حل على شكل سلسلة تامة حيث أن حساب حدودها سهل و بسيط كما أن الحلول التقريبية التي حصلنا عليها تنطبق على الحلول الحقيقية مما يؤكد أن هذه الطريقة أسهل و أيسر و أقوى من الطرق التقليدية مثل طرق: الفروق المنتهية، العناصر المنتهية، الطيفية..... إلخ

الكلمات المفتاحية: طريقة التفكيك، الشروط الحدية غير الموضعية، المعادلات التفاضلية ذات المشتقات الجزئية، الحل التحليلي.

I - INTRODUCTION

Partial differential equations with non local boundary conditions and partial integro-differential equations arise in many fields of science and engineering such as chemical diffusion, heat conduction processes, population dynamics, thermo-elasticity, medical science, electrochemistry and control theory [6-21]. A detailed description of the occurrence of such equations is given in [9]. In this paper, we consider a two-dimensional diffusion equation with non local boundary conditions. This type of problem was solved by many searchers using traditional numerical methods. For example, M. Siddique [1] proposed a fourth-order Padè-scheme. The purpose of this work is to study and use Adomian decomposition method [2-5] for this important problem. We show that this approach allows us to obtain an analytical solution. These results demonstrate that the decomposition method is more accurate, efficient and reliable in comparison with the conventional methods, like finite difference method etc...

We consider the two-dimensional diffusion equation given by

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2}, \quad 0 < x, y < 1, t > 0 \quad (1)$$

Initial conditions are assumed to be of the form

$$\begin{aligned} u(x, y, 0) &= f(x, y), \quad (x, y) \in \Omega \cup \partial\Omega \\ \text{And the Dirichelet time-dependent boundary conditions are:} \\ u(0, y, t) &= \psi_0(y, t), \quad 0 \leq t \leq T, 0 \leq y \leq 1 \\ u(1, y, t) &= \psi_1(y, t), \quad 0 \leq t \leq T, 0 \leq y \leq 1 \end{aligned} \quad (2)$$

$$\begin{aligned} u(x, 0, t) &= \varphi_0(x) \gamma(t), \quad 0 \leq t \leq T, 0 \leq x \leq 1 \\ u(x, 1, t) &= \varphi_1(x, t), \quad 0 \leq t \leq T, 0 \leq x \leq 1 \\ \text{And non local boundary condition:} \\ \int_0^1 \int_0^1 u(x, y, t) dx dy &= m(t), \quad (x, y) \in \Omega \cup \partial\Omega \end{aligned} \quad (3)$$

Where $f, \psi_0, \psi_1, \varphi_0, \varphi_1$ and m are known functions and $\gamma(t)$ is to be determined.

II adomian decomposition method

A. Operator form

In this section, we outline the steps to obtain a solution the above problem using the Adomian decomposition method, which was initiated by G.Adomian [9-11]. For this purpose, it is convenient to rewrite the problem in the standard form:

$$L_t(u(x, y, t)) = L_{xx}(u(x, y, t)) + L_{yy}(u(x, y, t)) \quad (4)$$

Where the differential operators $L_t(\cdot) = \frac{\partial}{\partial t}(\cdot)$, $L_{xx} = \frac{\partial^2}{\partial x^2}$ and $L_{yy} = \frac{\partial^2}{\partial y^2}$.

Assuming that the inverse L_t^{-1} exists and is defined as:

$$L_T^{-1} = \int_0^t (\cdot) dt \quad (5)$$

B. Application to the problem

Applying inverse operator on both the sides of (4) and using the initial condition, yields:

$$\begin{aligned} u(x, y, t) &= L_t^{-1} \left(L_{xx}(u(x, y, t)) + L_{yy}(u(x, y, t)) \right) \\ \text{or} \\ u(x, y, t) &= u(x, y, 0) + L_t^{-1} \left(L_{xx}(u(x, y, t)) + L_{yy}(u(x, y, t)) \right) \end{aligned} \quad (6)$$

Now, we decompose the unknown function $u(x,y,t)$ as a sum of components defined by the series :

$$u(x, y, t) = \sum_{k=0}^{\infty} u_k(x, y, t) \quad (7)$$

Where $u_0(x, y, t)$ is identified as $u(x, y, 0)$. Substituting equation (7) into equation (6) one obtains

$$\sum_{k=0}^{\infty} u_k(x, y, t) = f(x, y) + L_t^{-1} \{ L_{xx}(\sum_{k=0}^{\infty} u_k(x, y, t)) + L_{yy}(\sum_{k=0}^{\infty} u_k(x, y, t)) \} \quad (8)$$

The components $u_k(x, y, t)$ are obtained by the recursive formula:

$$u_0(x, y, t) = f(x, y) \quad (9)$$

$$u_{k+1}(x, y, t) = L_t^{-1} \left(L_{xx}(u_k(x, y, t)) + L_{yy}(u_k(x, y, t)) \right), \quad k \geq 0 \quad (10)$$

From equations (9) and (10) we obtain the first few terms as:

$$\begin{aligned} u_0 &= f(x, y) \\ u_1 &= L_t^{-1} (L_{xx}(u_0(x, y, t)) + L_{yy}(u_0(x, y, t))) \end{aligned}$$

$$u_2 = L_t^{-1} (L_{xx}(u_1(x, y, t)) + L_{yy}(u_1(x, y, t)))$$

$$u_3 = L_t^{-1} (L_{xx}(u_2(x, y, t)) + L_{yy}(u_2(x, y, t)))$$

and so on. As a result, the components u_0, u_1, u_2, \dots are identified and the series solution is thus entirely determined. However, in many cases the exact solution in a closed form may be obtained as we can see in our examples.

III. NUMERICAL EXAMPLES

A. Example 1

We consider the two-dimensional diffusion equation (1):

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2}$$

In which $u = u(x, y, t)$.

The Dirichelet time-dependent boundary conditions on the boundary $\partial\Omega$ of the square Ω defined by the line $x=0, y=0, x=1, y=1$ are given by:

$$u(x, 0, t) = e^{(x+2t)} \quad 0 \leq t \leq T, 0 \leq x \leq 1 \quad (11)$$

$$u(x, 1, t) = e^{(1+x+2t)} \quad 0 \leq t \leq T, \quad 0 \leq x \leq 1$$

And the non local boundary condition:

$$\int_0^1 \int_0^1 u(x, y, t) dx dy = (e - 1)^2 e^{2t} \quad (12)$$

With the initial conditions

$$u(x, y, 0) = e^{(x+y)} \quad (13)$$

Theoretical solution is given by :

$$u(x, y, t) = e^{(x+y+2t)} \quad (14)$$

Using the Adomians method, described above, equation (9) gives the first component

$$u_0(x, y, t) = f(x, y) = e^{(x+y)} \quad (15)$$

And equation (10) gives the following components of the series:

$$u_1 = L_t^{-1} (L_{xx}(u_0) + L_{yy}(u_0)) = 2 \int_0^t e^{x+y} dt = 2te^{x+y} \quad (16)$$

$$u_2 = L_t^{-1} (L_{xx}(u_1) + L_{yy}(u_1)) = 4 \int_0^t te^{x+y} dt = 2t^2 e^{x+y} \quad (17)$$

$$u_3 = L_t^{-1} (L_{xx}(u_2) + L_{yy}(u_2)) = 4 \int_0^t t^2 e^{x+y} dt = \frac{4}{3} t^3 e^{x+y} \quad (18)$$

$$u_4 = L_t^{-1} (L_{xx}(u_3) + L_{yy}(u_3)) = \frac{8}{3} \int_0^t t^3 e^{x+y} dt = \frac{2}{3} t^4 e^{x+y} \quad (19)$$

$$u_5 = L_t^{-1} (L_{xx}(u_4) + L_{yy}(u_4)) = \frac{4}{3} \int_0^t t^4 e^{x+y} dt = \frac{4}{3 \times 5} t^5 e^{x+y} \quad (20)$$

$$u_6 = L_t^{-1} (L_{xx}(u_5) + L_{yy}(u_5)) = \frac{8}{3 \times 5} \int_0^t t^5 e^{x+y} dt = \frac{8}{3 \times 5 \times 6} t^6 e^{x+y} \quad (21)$$

$$u_7 = L_t^{-1} (L_{xx}(u_6) + L_{yy}(u_6)) = \frac{16}{3 \times 5 \times 6} \int_0^t t^6 e^{x+y} dt = \frac{16}{3 \times 5 \times 6 \times 7} t^7 e^{x+y} \quad (22)$$

Substituting (15)-(22) into equation (7), we obtain the solution $u(x, y, t)$ of (1) with (11), and (12) in series form as:

$$u(x, y, t) = e^{x+y} \left(1 + \frac{2}{1!} t + \frac{4}{2!} t^2 + \frac{4 \times 2}{3!} t^3 + \frac{2 \times 2 \times 4}{4!} t^4 + \frac{2 \times 4 \times 4}{5!} t^5 + \frac{2 \times 4 \times 8}{6!} t^6 + \frac{2 \times 4 \times 16}{7!} t^7 + \dots \right) \quad (23)$$

Which can be rewritten as :

$$u(x, y, t) = e^{x+y} \left(1 + \frac{2t}{1!} + \frac{2^2 t^2}{2!} + \frac{2^3 t^3}{3!} + \frac{2^4 t^4}{4!} + \frac{2^5 t^5}{5!} + \frac{2^6 t^6}{6!} + \frac{2^7 t^7}{7!} + \dots \right) \quad (24)$$

It can be easily observed that (24) is equivalent to the exact solution:

$$u(x, y, t) = e^{(x+y)} e^t = e^{(x+y+2t)} \quad (25)$$

B. Example 2

Consider the two-dimensional nonhomogeneous diffusion problem

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} - e^{-t}(x^2 + y^2 + 4), \quad t > 0, x > 0, y < 1 \quad (26)$$

With Initial condition

$$u(x, y, 0) = 1 + x^2 + y^2 \quad (27)$$

And the boundary conditions:

$$\begin{aligned} u(0, y, t) &= 1 + y^2 e^{-t} \quad 0 \leq t \leq 1, 0 \leq y \leq 1 \\ u(1, y, t) &= 1 + (1 + y^2) e^{-t} \quad 0 \leq t \leq 1, 0 \leq y \leq 1 \end{aligned} \quad (28)$$

$$u(x, 0, t) = 1 + x^2 e^{-t}, \quad 0 \leq t \leq 1, 0 \leq x \leq 1$$

$$u(x, 1, t) = 1 + (1 + x^2) e^{-t}, \quad 0 \leq t \leq 1, 0 \leq x \leq 1$$

And the non local boundary condition

$$\int_0^1 \int_0^1 u(x, y, t) dx dy = 1 + \frac{2}{3} e^{-t}, \quad 0 \leq x \leq 1, 0 \leq y \leq 1 \quad (29)$$

The exact solution is:

$$u(x, y, t) = 1 + e^{-t}(x^2 + y^2) \quad (30)$$

Writing the problem in operator form and applying the inverse operator one obtains;

$$L_t^{-1} (L_t(u(x, y, t))) = L_t^{-1} (L_{xx}u(x, y, t)) + L_t^{-1} (L_{yy}u(x, y, t)) + L_t^{-1} (-e^{-t}(x^2 + y^2 + 4)) \quad (31)$$

$$L_t^{-1} (L_t(u(x, y, 0))) = u(x, y, 0) \quad (32)$$

From which we obtain:

$$u(x, y, t) = u(x, y, 0) + L_t^{-1} (L_{xx}u(x, y, t)) + L_t^{-1} (L_{yy}u(x, y, t)) + L_t^{-1} (-e^{-t}(x^2 + y^2 + 4)) \quad (33)$$

Using Adomian decomposition, the zeroth component is given by:

$$u_0(x, y, t) = u(x, y, 0) + L_t^{-1} (-e^{-t}(x^2 + y^2 + 4)) \quad (34)$$

and

$$u_{k+1}(x, t) = L_t^{-1} (L_{xx}(u_k(x, y, t)) + L_{yy}(u_k(x, y, t))), \quad k \geq 0 \quad (35)$$

Applying these formula, we obtain the components of the series as :

$$\begin{aligned} u_0(x, y, t) &= 1 + x^2 + y^2 - (x^2 + y^2 + 4) \int_0^t e^{-t} dt \\ u_0(x, y, t) &= -3 + (4 + x^2 + y^2)e^{-t} \end{aligned} \quad (36)$$

$$\begin{aligned} u_1(x, y, t) &= L_t^{-1} \left((L_{xx} + L_{yy})(u_0(x, y, t)) \right) = \int_0^t 4e^{-t} dt \\ u_1(x, y, t) &= -4e^{-t} + 4 \end{aligned} \quad (37)$$

$$\begin{aligned} u_2(x, y, t) &= L_t^{-1} \left((L_{xx} + L_{yy})(u_1(x, y, t)) \right) = \int_0^t 0 dt = 0 \\ u_k(x, y, t) &= 0, \quad k \geq 2 \end{aligned} \quad (38)$$

Once the components are determined then, the series solution completely determined as follows:

$$\begin{aligned} u(x, y, t) &= u_0(x, y, t) + u_1(x, y, t) + \sum_{k=2}^{\infty} u_k(x, y, t) \\ u(x, y, t) &= 1 + (x^2 + y^2)e^{-t} \end{aligned} \quad (39)$$

This solution coincides with the exact one.

C. Example 3.

Consider the two-dimensional diffusion problem (1)

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2}, \quad 0 < x, y < 1, t > 0$$

Subject to the initial condition

$$u(x, y, 0) = (1 - y)e^x, \quad 0 \leq x, y \leq 1 \quad (40)$$

And the boundary conditions

$$\begin{aligned} u(0, y, t) &= (1 - y)e^t, \quad 0 \leq t \leq 1, 0 \leq y \leq 1 \\ u(1, y, t) &= (1 - y)e^{1+t}, \quad 0 \leq t \leq 1, 0 \leq y \leq 1 \end{aligned} \quad (41)$$

$$\begin{aligned} u(x, 0, t) &= e^{1+t}, \quad 0 \leq t \leq 1, 0 \leq x \leq 1 \\ u(x, 1, t) &= 0, \quad 0 \leq t \leq 1, 0 \leq x \leq 1 \end{aligned}$$

Rewriting the equation (48) in an operator form as the follows

$$u_0(x, y, t) = u(x, y, 0) \quad (42)$$

$$u_{k+1}(x, y, t) = L_t^{-1} \left((L_{xx} + L_{yy})(u_k(x, y, t)) \right), \quad k \geq 0 \quad (43)$$

Using the Adomians method, described above, equation (9) gives the first component of the series as:

$$u_0(x, y, t) = (1 - y)e^x \quad (44)$$

And the remaining components are obtained using (10) which gives

$$u_1(x, y, t) = \int_0^t (1 - y)e^x dt = (1 - y)te^x \quad (45)$$

$$u_2(x, y, t) = \int_0^t (1 - y)te^x dt = (1 - y)e^x \left(\frac{t^2}{2!}\right) \quad (46)$$

$$u_3(x, y, t) = \int_0^t (1 - y)\left(\frac{t^2}{2!}\right)e^x dt = (1 - y)e^x \left(\frac{t^3}{3!}\right) \quad (47)$$

$$\dots\dots\dots u_k(x, y, t) = \int_0^t (1 - y)\left(\frac{t^{k-1}}{(k-1)!}\right)e^x dt = (1 - y)e^x \left(\frac{t^k}{k!}\right) \quad (48)$$

Using these results, the solution in series form is given by;

$$u(x, y, t) = (1 - y)e^x \left(\sum_{k=0}^{\infty} \frac{t^k}{k!}\right) = (1 - y)e^{x+t} \quad (49)$$

As we can verify by substitution, this solution is equivalent to the theoretical one.

CONCLUSION

In this work we applied the Adomian decomposition method for solving a two-dimensional diffusion equation, the method avoid the difficulties and massive computational work by determining the theoretical solution. The obtained solution is very rapidly convergent and provides a reliable technique that requires less effort and yields highly accurate results in comparison with the traditional techniques. The Adomian methodology is very powerful and efficient tool in finding exact solutions for a wide classes of problems. Through the illustrated problems, the obtained results are more accurate than those obtained by M. Siddique

REFERENCES

[1] M.Siddique, "Numerical Computatinal of Two-dimensional Diffusion Equation with Nonlocal Boundary Conditions," IAENG international journal of applied mathematics, 40:1, IJAM_40_1_04

[2] G.Adomian, *Solving Frontier Problems of Physics: The Decomposition Method*, Kluver Academic Publishers, Boston 1994.

[3] G.Adomian and R.Rach, "Noise Terms in Decomposition Solution series," Computers Math. Appl, 24(11) (1992),61-4.

[4] G. Adomian, "A review of The decomposition Method in Applied Mathematics," J.Math.Anal.Appl, 135 (1988),501-544.

[5] A.M.Wazwaz, "Necessary Conditions for The appearance of Noise Terms in Decomposition Solution Series," J.Math.Anal.Appl. 5 (1997),265-274.

[6] Cannon,J.R ,Y.Lin and S.Wang. "An Implicit Finite Difference Scheme for the Diffusion Equation Subject to Mass Specification", Int.J.Eng.Sci.28 (1990),573-578.

[7] Cannon;J.R and Van der Hoek, J. "Diffusion equation Subject to the Specification of Mass," J.Math.Anal.Appl., 115.pp.517-529,1986.

[8] Capsso,V. and Kunish, K.A, " reaction Diffusion stem Arising in Modeling Man-Environment Diseases," Q.Appl.Math.46, pp 431-439.1988

[9] Day.W.A, "Existence of A property of Solutions of The heat to Linear Thermoelasticity And other Theories," Quart.Appl.Math. 40, pp.319- 330,1982.

- [10] Wang.S. "A numerical Method for The heat Conduction Subject to Moving Boundary Energy Specification," Intern.J.Engng.Sci, 28, 543-546, 1991
- [11] Noye,B.J. and Hayman, K.J.Explicit, "Two Level Finite Difference Methods for the Two-Dimensional Diffusion Equation," Intern. J. Computr. Math.42, pp.223-236.1992
- [12] Wang.S. and Lin.Y.,A Finite Difference Solution to An Inverse Problem Determining a Controlling Function in a Parabolic Partial Differential Equation. Inverse Problems.5,pp.631-640.
- [13] B.A.Wade, A.Q.M.Khaliq ,M.Siddique and M.Yousuf, "Smoothing with Positivity-Preserving Padè Schemes for Parabolic Problems with Non Smooth Data", Numerical Methods for Partial Differential Equations (NMPDE),Wiley Interscience;V.21.No 3.2005,pp.553- 573.
- [14] B.A.Wade,A.Q.M.Khaliq,M.Yousuf and J.Vigo-Aguiar" High-Order Smoothing Schemes For Inhomogeneous Parabolic problems with Application to Non smooth Payoff In option Pricing" Numerical Methods for Partial Differential Equations (NMPDE) V.23(5).2007,1249-1276. 10
- [15] A.B.Gumel,W.T.Ang And F.H.Twizell."Efficient Parallel Algorithm for The Two-Dimensional Diffusion Equation Subject to Specification of Mass" Intern. J. Computer Math. Vol 64,p 153-163 (1997).
- [16] G.D.Smith, "Numerical Solution for Partial Differential Equations Finite Difference Methods", Third Edition, Oxford University Press, New york (1985).
- [17] A.Q.M.Khaliq,E.H.Twizell And D.A.Voss."On Parallel Algorithm for Sem-discretized Parabolic Partial Differential Equations Based On Subdiagonal Padè Approximation" (NMPDE),Wiley Interscience.9.107-116 (1993).
- [18] P.Marcati,Some Considerations ON the Mathematical Approach to Nonlinear Age Dependent Population Dynamics, Computers Math.Applie.9 (3) 361-370 (1983).
- [19] Y.Lin And S.Wang."A Numerical Method for the Diffusion Equation with Nonlocal Boundary Conditions"Int.J.Eng.Sci.28 (1990),543-546.
- [20] M.Siddique."A Comparison of Third-Order the Diffusion Equation L0-Stable numerical Schemes for the Two-Dimensional Homogeneous Diffusion Problem Subject to Specification of Mass" Applied Mathematical Sciences.Vol.4.2010.No.13.611-621.
- [21] M.Siddique."Smoothing of Crank-Nicolson Schemes for the Two-Dimensional Diffusion with an Integral Condition" Applied Mathematics and Computation. 214.2009.512-522.